

Glucosamine Plus™ Extra Strength

DESCRIPTION

Glucosamine Plus Extra Strength capsules, provided by Douglas Laboratories®, contain a synergistic combination of glucosamine sulfate and chondroitin sulfate. Glucosamine sulfate is an important precursor for the synthesis and maintenance of connective tissues. Chondroitin sulfate also supports formation of connective tissues. The potassium stabilized form of glucosamine sulfate is used, making this sodium-free product.

FUNCTIONS

Glucosamine is a naturally occurring amino sugar found ubiquitously in glycoproteins and glycosaminoglycans. Glycosaminoglycans, formerly named mucopolysaccharides, are an integral component of all connective tissues. Connective tissue, a fibrous type of body tissue, has various functions. It supports and connects internal organs (ligaments), forms bone, cartilage, and the walls of blood vessels, attaches muscles to bones (tendons), and replaces tissues that have been damaged following injury.

The two main components of connective tissue are collagen and proteoglycans. Collagen is the strong, fibrous protein that physically connects our tissues. Proteoglycans are large carbohydrate-rich structures, resembling a bottlebrush in three-dimensional structure, i.e., a central protein rod with many strings of glucosamine-containing glycosaminoglycans extending outwards. Proteoglycans hold large amounts of water forming a stiff gel by virtue of their dense negative charges from sulfates. Proteoglycans are linked to collagen fibers to help form connective tissues, and proteoglycans provide resiliency, load distribution, shock-absorbing, compressive and lubricating properties to connective tissues and joints.

Glycosaminoglycans and proteoglycans are continuously being formed and replaced in connective tissues. Remodeling of connective tissues is also continuous, albeit slowly, with turnover half-lives of almost two years in healthy human joints. Turnover is accelerated in wound healing, arthritic joints, and burns. New synthesis of glycosaminoglycans has clinical importance in skin during normal wound healing, in bone during fracture repair and osteoporosis, and in joints. In these instances, glycosaminoglycan synthesis is necessary for healing, and enhancement of glycosaminoglycan and proteoglycan deposition may improve tissue repair. Glycosaminoglycans are synthesized primarily by fibroblasts (skin, tendons, ligaments), osteoblasts (bone), and chondrocytes (cartilage). Thus, the cell's ability to manufacture glycosaminoglycans and secrete proteoglycans is crucial during any healing or joint disease process.

The availability of glucosamine is the key, rate-limiting step in glycosaminoglycan and proteoglycan synthesis in all connective tissues, such as skin, bone, cartilage, tendons, and ligaments. Only with sufficient glucosamine, the synthesis can proceed.

The body has a long metabolic pathway to synthesize a glucosamine derivative, UDP-N-acetylglucosamine, which is used for glycosaminoglycan synthesis. In addition, exogenous, i.e., dietary, glucosamine serves as an immediate precursor for glycosaminoglycan synthesis, and also stimulates incorporation of other precursors into the connective tissue matrix.

Chondroitin sulfate, a glycosaminoglycan formed in the body, is also used for the synthesis and maintenance of connective tissue, primarily within the cartilage matrix. In addition, chondroitin sulfate protects existing cartilage by reducing water loss from the matrix and by inhibiting the enzymatic breakdown of the cartilage. The overlapping activities as well as functional differences of glucosamine and chondroitin sulfate offer several advantages for combined supplementation. Although glucosamine is a precursor for chondroitin synthesis, this process requires large amounts of metabolic energy. Dietary preformed chondroitin sulfate spares the use of glucosamine for this purpose. Instead glucosamine can be used for formation of other important glycosaminoglycans and proteoglycans. When adequate chondroitin sulfate is thus available to help protect tissues from premature breakdown, glucosamine can more readily stimulate synthesis of healthy new tissue. Glucosamine is almost universally found in small amounts in most foods.

Glucosamine Plus™ Extra Strength

However, bioavailability of glucosamine from foods is largely unknown.

Bioavailability of oral glucosamine sulfate is excellent. It is absorbed intact, and utilized very quickly by all tissues, including connective tissues. Chondroitin sulfate is found in most animal tissues. Orally ingested chondroitin sulfate is also well absorbed and distributed to tissues.

In summary, glucosamine sulfate is a well documented, highly effective source of glucosamine for glycosaminoglycan and proteoglycan synthesis in all connective tissues. Chondroitin sulfate is an excellent source of n-acetylgalactosaminoglycan for synthesis and protection of proteoglycans associated with cartilage tissues.

INDICATIONS

Glucosamine Plus Extra Strength capsules may be a useful nutritional adjunct for individuals who wish to support the body's connective tissues, such as skin, tendons, ligaments, bone, and cartilage.

FORMULA (GSXV)

One Vegetarian Capsule Contains:

Glucosamine Sulfate.2KCl	500 mg
(from crab and shrimp)	
Chondroitin Sulfate.....	400 mg

SUGGESTED USE

Adults take one vegetarian capsule up to three times daily or as directed by a healthcare professional.

SIDE EFFECTS

No adverse side effects have been reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

Warning: If you have an allergy to shellfish, (including crab and shrimp) you should not use this product.

Glucosamine Plus™ Extra Strength

REFERENCES

- Bagchi M, Balmoori J, Bagchi D, Ray SD, Kuszynski C, Stohs SJ. Smokeless tobacco, oxidative stress, apoptosis, and antioxidants in human oral keratinocytes [published erratum appears in Free Radic Biol Med 1999 Jun;26(11-12):1599]. Free Radic Biol Med 1999;26:992-1000.
- Cranton EM, Frackleton JP: Free radical pathology in age-associated disease: treatment with EDTA chelation, nutrition, and antioxidants. J Holistic Med 1984;6:1-36.
- Davis BP et al: Enhanced absorption of oral vitamin B12 from a resin ascorbate administered to normal subjects. Manip Physiol Ter 1982;5:123-127.
- Feher J, Lengyel G, Blazovics A. Oxidative stress in the liver and biliary tract diseases. Scand J Gastroenterol Suppl 1998;228:38-46.
- Forsleff L, Schauss AG, Bier ID, Stuart S. Evidence of functional zinc deficiency in Parkinson's disease. J Altern Complement Med 1999;5:57-64.
- Johnson K, Kligman EW. Preventive nutrition: an 'optimal' diet for older adults. Geriatrics 1992;47:56-60.
- Oldham KM, Bowen PE. Oxidative stress in critical care: is antioxidant supplementation beneficial? J Am Diet Assoc 1998;98:1001-8.
- Seekamp A, Hultquist DE, Till GO. Protection by vitamin B2 against oxidant-mediated acute lung injury. Inflammation 1999;23:449-60.

For more information visit douglaslabs.com

† These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by
Douglas Laboratories
600 Boyce Road
Pittsburgh, PA 15205
800-245-4440
douglaslabs.com



**You trust Douglas Laboratories.
Your patients trust you.**